

ON A NEW CLASS OF RIGID COXETER GROUPS

TETSUYA HOSAKA

ABSTRACT. In this paper, we give a new class of rigid Coxeter groups. Let (W, S) be a Coxeter system. Suppose that (0) for each $s, t \in S$ such that $m(s, t)$ is even, $m(s, t) \in \{2\} \cup 4\mathbb{N}$, (1) for each $s \neq t \in S$ such that $m(s, t)$ is odd, $\{s, t\}$ is a maximal spherical subset of S , (2) there does not exist a three-points subset $\{s, t, u\} \subset S$ such that $m(s, t)$ and $m(t, u)$ are odd, and (3) for each $s \neq t \in S$ such that $m(s, t)$ is odd, the number of maximal spherical subsets of S intersecting with $\{s, t\}$ is at most two, where $m(s, t)$ is the order of st in the Coxeter group W . Then we show that the Coxeter group W is rigid. This is an extension of a result of D. Radcliffe.

1. INTRODUCTION AND PRELIMINARIES

The purpose of this paper is to give a new class of rigid Coxeter groups. A *Coxeter group* is a group W having a presentation

$$\langle S \mid (st)^{m(s,t)} = 1 \text{ for } s, t \in S \rangle,$$

where S is a finite set and $m : S \times S \rightarrow \mathbb{N} \cup \{\infty\}$ is a function satisfying the following conditions:

- (i) $m(s, t) = m(t, s)$ for any $s, t \in S$,
- (ii) $m(s, s) = 1$ for any $s \in S$, and
- (iii) $m(s, t) \geq 2$ for any $s, t \in S$ such that $s \neq t$.

The pair (W, S) is called a *Coxeter system*. For a Coxeter group W , a generating set S' of W is called a *Coxeter generating set for W* if (W, S') is a Coxeter system. Let (W, S) be a Coxeter system. For a subset $T \subset S$, W_T is defined as the subgroup of W generated by T , and called a *parabolic subgroup*. A subset $T \subset S$ is called a *spherical subset of S* , if the parabolic subgroup W_T is finite.

Date: January 5, 2005.

2000 Mathematics Subject Classification. 20F65, 20F55.

Key words and phrases. rigidity of Coxeter groups.

Partly supported by the Grant-in-Aid for Scientific Research, The Ministry of Education, Culture, Sports, Science and Technology, Japan, (No. 15740029).

Let (W, S) and (W', S') be Coxeter systems. Two Coxeter systems (W, S) and (W', S') are said to be *isomorphic*, if there exists a bijection $\psi : S \rightarrow S'$ such that

$$m(s, t) = m'(\psi(s), \psi(t))$$

for every $s, t \in S$, where $m(s, t)$ and $m'(s', t')$ are the orders of st in W and $s't'$ in W' , respectively.

In general, a Coxeter group does not always determine its Coxeter system up to isomorphism. Indeed some counter-examples are known (cf. [1], [2]). Here there exists the following natural problem.

Problem ([2], [3]). When does a Coxeter group determine its Coxeter system up to isomorphism?

A Coxeter group W is said to be *rigid*, if the Coxeter group W determines its Coxeter system up to isomorphism (i.e., for each Coxeter generating sets S and S' for W the Coxeter systems (W, S) and (W, S') are isomorphic).

A Coxeter system (W, S) is said to be *even*, if $m(s, t)$ is even for all $s \neq t$ in S . Also a Coxeter system (W, S) is said to be *strong even*, if $m(s, t) \in \{2\} \cup 4\mathbb{N}$ for all $s \neq t$ in S .

The following theorem was proved by D. Radcliffe in [5].

Theorem 1.1 ([5]). *If (W, S) is a strong even Coxeter system, then the Coxeter group W is rigid.*

In this paper, we prove the following theorem which is an extension of Theorem 1.1 and [4, Theorem 1.2].

Theorem 1.2. *Let (W, S) be a Coxeter system. Suppose that*

- (0) *for each $s, t \in S$ such that $m(s, t)$ is even, $m(s, t) \in \{2\} \cup 4\mathbb{N}$,*
- (1) *for each $s \neq t \in S$ such that $m(s, t)$ is odd, $\{s, t\}$ is a maximal spherical subset of S ,*
- (2) *there does not exist a three-points subset $\{s, t, u\} \subset S$ such that $m(s, t)$ and $m(t, u)$ are odd, and*
- (3) *for each $s \neq t \in S$ such that $m(s, t)$ is odd, the number of maximal spherical subsets of S intersecting with $\{s, t\}$ is at most two.*

Then the Coxeter group W is rigid.

2. PROOF OF THE THEOREM

Let (W, S) be a Coxeter system. Suppose that

- (0) for each $s, t \in S$ such that $m(s, t)$ is even, $m(s, t) \in \{2\} \cup 4\mathbb{N}$,

- (1) for each $s \neq t \in S$ such that $m(s, t)$ is odd, $\{s, t\}$ is a maximal spherical subset of S ,
- (2) there does not exist a three-points subset $\{s, t, u\} \subset S$ such that $m(s, t)$ and $m(t, u)$ are odd, and
- (3) for each $s \neq t \in S$ such that $m(s, t)$ is odd, the number of maximal spherical subsets of S intersecting with $\{s, t\}$ is at most two.

Let (W', S') be a Coxeter system. We suppose that there exists an isomorphism $\phi : W \rightarrow W'$. To prove Theorem 1.2, we show that the Coxeter systems (W, S) and (W', S') are isomorphic.

The following lemma is known.

Lemma 2.1 (cf. [2], [5]). *For each maximal spherical subset $T \subset S$, there exists a unique maximal spherical subset $T' \subset S'$ such that $\phi(W_T) = w'W'_{T'}w'^{-1}$ for some $w' \in W'$.*

We first prove the following lemma.

Lemma 2.2. *The Coxeter system (W', S') satisfies the following:*

- (0') for each $s', t' \in S'$ such that $m'(s', t')$ is even, $m'(s', t') \in \{2\} \cup 4\mathbb{N}$,
- (1') for each $s' \neq t' \in S'$ such that $m'(s', t')$ is odd, $\{s', t'\}$ is a maximal spherical subset of S' ,
- (2') there does not exist a three-points subset $\{s', t', u'\} \subset S'$ such that $m'(s', t')$ and $m'(t', u')$ are odd, and
- (3') for each $s' \neq t' \in S'$ such that $m'(s', t')$ is odd, the number of maximal spherical subsets of S' intersecting with $\{s', t'\}$ is at most two.

Proof. Let $s' \neq t' \in S'$. There exists a maximal spherical subset T' of S' such that $\{s', t'\} \subset T'$. By Lemma 2.1, $\phi^{-1}(W'_{T'}) \sim W_T$ for some maximal spherical subset T of S . By (0) and (1), either

- (i) (W_T, T) is a strong even Coxeter system, or
- (ii) $|T| = 2$ and if $T = \{s, t\}$ then $m(s, t)$ is odd.

Hence W_T is a rigid Coxeter group and (W_T, T) and $(W'_{T'}, T')$ are isomorphic. Thus if $m'(s', t')$ is even then $m'(s', t') \in \{2\} \cup 4\mathbb{N}$, and if $m'(s', t')$ is odd then $\{s', t'\}$ is a maximal spherical subset of S' . Hence (0') and (1') hold. We can show (2') and (3') by the same argument as the proof of [4, Lemma 3.1] $\square$

Let $\mathcal{A}$ and $\mathcal{A}'$ be the sets of all maximal spherical subsets of S and S' , respectively. For each $T \in \mathcal{A}$, there exists a unique element $T' \in \mathcal{A}'$ such that $\phi(W_T) \sim W'_{T'}$ by Lemma 2.1.

We define

$$\begin{aligned}\bar{S} &= \bigcup \{T \in \mathcal{A} \mid (W_T, T) \text{ is strong even}\} \\ \bar{S}' &= \bigcup \{T' \in \mathcal{A}' \mid (W'_{T'}, T') \text{ is strong even}\}.\end{aligned}$$

We note that for each $s \in S \setminus \bar{S}$, there exists a unique element $t \in S \setminus \{s\}$ such that $m(s, t)$ is odd. Then $m(s, u) = \infty$ for any $u \in S \setminus \{s, t\}$.

Let W^{ab} and W'^{ab} be the abelianizations of W and W' respectively, and let $\pi : W \rightarrow W^{\text{ab}}$ and $\pi' : W' \rightarrow W'^{\text{ab}}$ be the abelianization maps. We note that $W^{\text{ab}} = (W_{\bar{S}})^{\text{ab}} \cong \mathbb{Z}_2^{|\bar{S}|}$ and $W'^{\text{ab}} = (W'_{\bar{S}'})^{\text{ab}} \cong \mathbb{Z}_2^{|\bar{S}'|}$.

We can obtain the following lemma by the same argument as the proof of [5, Theorem 4.4].

Lemma 2.3. *If A and B are subsets of $\bar{S}$ and $\pi(W_A) = \pi(W_B)$, then $A = B$.*

For $A \subset \bar{S}$ and $A' \subset \bar{S}'$, we denote $A\tau A'$ if $\pi'(\phi(W_A)) = \pi'(W'_{A'})$.

We can obtain the following lemma by the same argument as the proof of [5, Theorem 4.5].

Lemma 2.4. *Let A and B be subsets of $\bar{S}$ and let A' and B' be subsets of $\bar{S}'$.*

- (i) *If $A\tau A'$ and $B\tau A'$ then $A = B$.*
- (ii) *If $A\tau A'$ and $A\tau B'$ then $A' = B'$.*
- (iii) *If $A\tau A'$ and $B\tau B'$ then $(A \cap B)\tau(A' \cap B')$.*

We obtain the following lemma from Lemmas 2.3 and 2.4.

Lemma 2.5. *Let A and B be subsets of $\bar{S}$ and let A' and B' be subsets of $\bar{S}'$. If $A\tau A'$, $B\tau B'$ and $A \subset B$, then $A' \subset B'$.*

Proof. Suppose that $A\tau A'$, $B\tau B'$ and $A \subset B$. By Lemma 2.4 (iii), $(A \cap B)\tau(A' \cap B')$. Since $A \subset B$, $A\tau(A' \cap B')$. Now $A\tau A'$. By Lemma 2.4 (ii), $A' = A' \cap B'$, i.e., $A' \subset B'$. $\square$

A subset T of S is said to be *independent*, if $m(s, t) = 2$ for all $s \neq t$ in T . We note that if T is an independent subset of S then $W_T \cong \mathbb{Z}_2^{|T|}$. Let $\mathcal{B}$ and $\mathcal{B}'$ be the sets of all maximal independent subsets of $\bar{S}$ and $\bar{S}'$, respectively.

We show the following lemma which correspond to [5, Theorem 4.7].

Lemma 2.6. *For each $T \in \mathcal{B}$, there exists a unique $T' \in \mathcal{B}'$ such that $T\tau T'$.*

Proof. Let $T \in \mathcal{B}$. Then there exists $U \in \mathcal{A}$ such that $T \subset U \subset \bar{S}$. By Lemma 2.1, $\phi(W_U) = w'W'_{U'}w'^{-1}$ for some $U' \in \mathcal{A}'$ and $w' \in W'$. Here $\phi : W_U \rightarrow w'W'_{U'}w'^{-1}$ is an isomorphism and (W_U, U) and $(W'_{U'}, U')$ are strong even. By the proof of [5, Theorem 4.7], there exists a unique independent subset T' of U' such that $T\tau T'$. We show that T' is a *maximal* independent subset of $\bar{S}'$. Suppose that $T' \subset T'_0$ and T'_0 is an independent subset of $\bar{S}$. Then by the above argument, there exists an independent subset T_0 of $\bar{S}$ such that $T_0\tau T'_0$. Since $T' \subset T'_0$, $T \subset T_0$ by Lemma 2.5. Hence $T = T_0$ because T is a maximal independent subset of $\bar{S}$. By Lemma 2.4 (ii), $T' = T'_0$. Thus T' is a maximal independent subset of $\bar{S}'$, i.e., $T' \in \mathcal{B}'$ which is a unique element such that $T\tau T'$. $\square$

We can obtain the following lemma from Lemmas 2.4 (iii) and 2.6 and the proof of [5, Theorem 4.8].

Lemma 2.7. *Let $T_1, \dots, T_k \in \mathcal{A} \cup \mathcal{B}$ and $T'_1, \dots, T'_k \in \mathcal{A}' \cup \mathcal{B}'$ such that $T_i \subset \bar{S}$ and $T_i\tau T'_i$ for each $i = 1, \dots, k$. Then $|T_1 \cap \dots \cap T_k| = |T'_1 \cap \dots \cap T'_k|$.*

Lemma 2.7 implies that there exists a bijection $\bar{\psi} : \bar{S} \rightarrow \bar{S}'$ such that for each $s \in \bar{S}$ and $T \in \mathcal{A} \cup \mathcal{B}$ with $T \subset \bar{S}$, $s \in T$ if and only if $\bar{\psi}(s) \in T'$, where T' is the element of $\mathcal{A}' \cup \mathcal{B}'$ such that $T\tau T'$ (cf. [5]). By the proof of [5, Theorem 4.11], the bijection $\bar{\psi} : \bar{S} \rightarrow \bar{S}'$ induces an isomorphism between the Coxeter systems $(W_{\bar{S}}, \bar{S})$ and $(W_{\bar{S}'}, \bar{S}')$.

Here we note that we can construct $\bar{\psi} : \bar{S} \rightarrow \bar{S}'$ so that $\bar{\psi}(t) = t'$ for each $t \in \bar{S}$ and $t' \in \bar{S}'$ such that $\{t\}\tau\{t'\}$. Indeed, suppose that $\{t\}\tau\{t'\}$ (such t' is unique, since $(W_{\bar{S}}, \bar{S})$ and $(W'_{\bar{S}'}, \bar{S}')$ are even). Then for $T \in \mathcal{A} \cup \mathcal{B}$ with $T \subset \bar{S}$ and $T' \in \mathcal{A}' \cup \mathcal{B}'$ such that $T\tau T'$, $t \in T$ if and only if $t' \in T'$ by Lemma 2.5.

Using the above argument, we show the following.

Theorem 2.8. *The Coxeter systems (W, S) and (W', S') are isomorphic.*

Proof. We define a bijection $\psi : S \rightarrow S'$ as follows: Let $s \in S$. If $s \in \bar{S}$ then we define $\psi(s) = \bar{\psi}(s)$. Suppose that $s \in S \setminus \bar{S}$. Then there exists a unique element $t \in S \setminus \{s\}$ such that $m(s, t)$ is odd. Here we note that $m(s, u) = \infty$ for any $u \in S \setminus \{s, t\}$. Now either $t \in \bar{S}$ or $t \notin \bar{S}$. We first suppose that $t \notin \bar{S}$, i.e., $\{s, t\} \subset S \setminus \bar{S}$. Then $\{T \in \mathcal{A} \mid T \cap \{s, t\} \neq \emptyset\} = \{\{s, t\}\}$. There exists a unique $\{s', t'\} \in \mathcal{A}'$ such that $\phi(W_{\{s, t\}}) \sim W'_{\{s', t'\}}$ by Lemma 2.1. Here $\{s', t'\} \subset S \setminus \bar{S}$ by [4, Lemma 2.6]. We define $\psi(s) = s'$ and $\psi(t) = t'$. Next we suppose that $t \in \bar{S}$. Then $|\{T \in \mathcal{A} \mid T \cap \{s, t\} \neq \emptyset\}| = 2$, and there exists a

unique $T \in \mathcal{A}$ such that $t \in T \subset \bar{S}$. By Lemma 2.1, there exist unique $\{s', t'\}, T' \in \mathcal{A}'$ such that $\phi(W_{\{s, t\}}) \sim W'_{\{s', t'\}}$ and $\phi(W_T) \sim W'_{T'}$. The proof of [4, Lemma 2.6] implies that $\{s', t'\} \cap T' \neq \emptyset$ and $\phi(t) \sim s' \sim t'$. We may suppose that $t' \in T'$. Then $\bar{\psi}(t) = t'$ because $\{t\}\tau\{t'\}$. We define $\psi(s) = s'$.

Then the bijection $\psi : S \rightarrow S'$ induces an isomorphism between the Coxeter systems (W, S) and (W', S') by the construction of ψ . $\square$

REFERENCES

- [1] N. Bourbaki, *Groupes et Algèbres de Lie*, Chapters IV-VI, Masson, Paris, 1981.
- [2] N. Brady, J.P. McCammond, B. Mühlherr and W.D. Neumann, *Rigidity of Coxeter groups and Artin groups*, *Geom. Dedicata* **94** (2002), 91–109.
- [3] R. Charney and M.W. Davis, *When is a Coxeter system determined by its Coxeter group?* *J. London Math. Soc.* **61** (no.2) (2000), 441–461.
- [4] T. Hosaka, *A class of rigid Coxeter groups*, preprint.
- [5] D. Radcliffe, *Unique presentation of Coxeter groups and related groups*, Ph.D. thesis, The University of Wisconsin-Milwaukee, 2001.

DEPARTMENT OF MATHEMATICS, UTSUNOMIYA UNIVERSITY, UTSUNOMIYA,
321-8505, JAPAN

E-mail address: hosaka@cc.utsunomiya-u.ac.jp